Medications at Time of Surgery

SPECIAL INFORMATION FOR PATIENTS OF

ATUL F. KAMATH, MD

▶ MEDICATION CHECKLIST

Prior to joint replacement surgery, it is very important that you stop taking certain types of medications and supplements. Please stop taking the following items at the listed timeframe prior to your surgery:

*NOTE: If you are taking DMARDs or other biologics, please consult with your rheumatologist regarding when to stop medication.

Seven Days Before Surgery:

- STOP Anti-coagulants / Anti-platelets / Anti-thrombotics (medications that prevent clotting)
- STOP Aspirin
- STOP Herbal supplements
- STOP Over-the-counter medications
- STOP Vitamins

*NOTE: You should stop hormonal therapy a week before surgery including birth control, tamoxifen (following discussion with oncologist) and estrogen replacement therapy.

Five Days Before Surgery:

- STOP NSAIDs (including Ibuprofen, Advil, Aleve and Motrin)
- · START Bactroban ointment, if prescribed

Two Days Before Surgery:

• START Sennakot (Stool Softener)

Morning of Surgery:

- STOP ACE-Inhibitors (medications for high blood pressure and congestive heart failure)
- STOP ARBs (medications for high blood pressure, kidney damage and congestive heart failure)

You should continue to take certain medications with a sip of water on the morning of your surgery including:

- · Beta blockers (including Toprol, Corgard, Levatol and Coreg)
- Long acting pain medications (including Oxycontin and MS Contin)
- Tylenol
- Celebrex

▶ DIABETIC MEDICATION

If you are diabetic, please take your medication as listed below:

Night before Surgery

• Take oral medication or insulin after last meal, as normal

Morning of Surgery

- Do not take oral medication
- · If taking insulin, contact your prescribing physician for advice on altered dosage

If you have any questions regarding your specific medications or medications that are not listed above, please consult you surgeon.



